



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

another instance of the specialization of the Swifts along various lines, and also as showing the structural variation among different members of the group. In *Macropteryx* the deltoid is well-developed and arises from two heads as in the Passeres. In *Cypseloides*, *Micropus*, *Collocalia*, and *Tachornis*, the deltoid is single and reduced in size, being proportionally smallest in *Tachornis*. Finally, as stated above, the deltoid is quite absent in *Chætura pelagica*, this being a step beyond what is found among the Hummingbirds, where the deltoid is present though small. It would be interesting to know if other members of the genus *Chætura* lack the deltoid, and also what is the condition of this muscle in *Hemiprocne*.

Dr. Shufeldt's figure of the wing muscles of *Chætura* (Linn. Soc. Journ. Zool. XX, pl. 22, fig. 29) shows that he too found this muscle absent, although nothing is said about it in the text.

I would be extremely grateful to any one for alcoholic specimens of *Hemiprocne* or any species of *Chætura* save *pelagica*.—F. A. LUCAS, Washington, D. C.

The Scissor-tailed Flycatcher in Virginia and Maryland.—My friend, Mr. P. Henry Azlett, of Azlett, King William County, Virginia, recently sent me for identification an adult specimen of this bird (*Milvulus forficatus*) which was shot by a farmer near that place on August 31, 1895. The bird is in poor, half moulted condition. This is, I believe, the fourth record of this bird for Virginia, and it is of course possible that some or all were escaped cage birds. The late Mr. O. N. Bryan of Bryan's Point, Maryland, on the Potomac River just below Washington, once told me that in August of a year about the close of the war while he was in a deep ravine near his home, called Johnson's Gully, he was overtaken by a severe storm, and saw one of these birds which had evidently sought the seclusion of the same place for shelter.—WILLIAM PALMER, Washington, D. C.

The Raven in Illinois.—I wish to place on record the capture of a specimen of *Corvus corax principalis*, at Mendosa, Ill., Oct. 23, 1892. I was at that time making a collection of birds for the State, and was living with the crew of the United States Fish Commission. The bird had been seen for a week or more previous to this time, flying about Lake Mendosa, a body of water opening into the Illinois River. The lake is seven miles long and three quarters of a mile wide. A number of attempts had been made by market shooters to obtain the bird for me, but they could not get within gunshot, and so were unsuccessful.

One day, however, as I was passing through a herd of cattle, the bird flew very low and I obtained it with a charge of number five shot. The specimen, a female, had been feeding upon carrion, and the odor from the body was as disagreeable as that from *Cathartes aura*. The plumage is exceptionally fine; the body is deep blue black; from the secondaries to the primary coverts, the color is rich brownish bronze. The following